

CP Violation and Strategies for Extracting CKM Phases

Robert Fleischer

Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Abstract

The phenomenon of CP violation and strategies for extracting the angles of the unitarity triangle are reviewed. Special emphasis is given to the B -meson system, as it plays an outstanding role to test the Standard Model description of CP violation at future experimental B -physics facilities. Both general aspects and recent developments are discussed. It is pointed out that combined branching ratios for $B_{u,d} \rightarrow \pi K$ modes, which have been observed recently by the CLEO collaboration, and untagged $B_s \rightarrow K\bar{K}$ decays may allow us to derive interesting constraints on the CKM angle γ and may eventually lead to a solid determination of this angle.

*Invited plenary talk given at the
IVth International Workshop on Progress in Heavy Quark Physics
Rostock, Germany, 20–22 September 1997
To appear in the Proceedings*

CP VIOLATION AND STRATEGIES FOR EXTRACTING CKM PHASES

ROBERT FLEISCHER
Theory Division, CERN
CH-1211 Geneva 23, Switzerland

ABSTRACT

The phenomenon of CP violation and strategies for extracting the angles of the unitarity triangle are reviewed. Special emphasis is given to the B -meson system, as it plays an outstanding role to test the Standard Model description of CP violation at future experimental B -physics facilities. Both general aspects and recent developments are discussed. It is pointed out that combined branching ratios for $B_{u,d} \rightarrow \pi K$ modes, which have been observed recently by the CLEO collaboration, and untagged $B_s \rightarrow K\bar{K}$ decays may allow us to derive interesting constraints on the CKM angle γ and may eventually lead to a solid determination of this angle.

1. Introduction

CP violation plays a central and fundamental role in present particle physics. Since it is one of the least experimentally explored phenomena of the Standard Model, it may still shed light on new physics. To this end, it is crucial to search for CP-violating processes that can be analysed in a clean way within the Standard Model framework.

Within the Standard Model of electroweak interactions, CP violation is closely related to the Cabibbo–Kobayashi–Maskawa matrix¹ (CKM matrix) connecting the electroweak eigenstates of the d , s and b quarks with their mass eigenstates. As far as CP violation is concerned, the central feature of the CKM matrix is that – in addition to three generalized Cabibbo-type angles – also a *complex phase* is needed in the three-generation case to parametrize the CKM matrix. This complex phase is the origin of CP violation within the Standard Model.

A closer look shows that CP-violating observables are proportional to the following combination of CKM matrix elements:

$$J_{\text{CP}} = \pm \text{Im} \left(V_{i\alpha} V_{j\beta} V_{i\beta}^* V_{j\alpha}^* \right) \quad (i \neq j, \alpha \neq \beta), \quad (1)$$

which represents a measure of the “strength” of CP violation within the Standard Model². Since $J_{\text{CP}} = \mathcal{O}(10^{-5})$, CP violation is a small effect. Typically several new complex couplings are present in many scenarios of new physics³, yielding additional sources for CP violation.

Concerning phenomenological applications, the parametrization

$$\hat{V}_{\text{CKM}} = \begin{pmatrix} 1 - \frac{1}{2}\lambda^2 & \lambda & A\lambda^3 R_b e^{-i\gamma} \\ -\lambda & 1 - \frac{1}{2}\lambda^2 & A\lambda^2 \\ A\lambda^3 R_t e^{-i\beta} & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4) \quad (2)$$

with $\lambda = 0.22$ and

$$A \equiv \frac{1}{\lambda^2} |V_{cb}| = 0.81 \pm 0.06 \quad (3)$$

$$R_b \equiv \frac{1}{\lambda} \left| \frac{V_{ub}}{V_{cb}} \right| = \sqrt{\rho^2 + \eta^2} = 0.36 \pm 0.08, \quad R_t \equiv \frac{1}{\lambda} \left| \frac{V_{td}}{V_{cb}} \right| = \sqrt{(1 - \rho)^2 + \eta^2} = \mathcal{O}(1) \quad (4)$$

turns out to be very useful. This parametrization is a modification of the Wolfenstein parametrization⁴ expliciting not only the hierarchy of the CKM elements, but also the dependence on the angles $\beta = \beta(\rho, \eta)$ and $\gamma = \gamma(\rho, \eta)$ of the usual “non-squashed” unitarity triangle of the CKM matrix⁵.

2. A Brief Look at CP Violation in Kaon Decays

So far, CP violation has been observed only within the neutral K -meson system, although the discovery of this phenomenon goes back to the year 1964⁶. In the neutral kaon system, CP violation is described by two complex quantities, called ε and ε' , which are defined by the following ratios of decay amplitudes:

$$\frac{A(K_L \rightarrow \pi^+ \pi^-)}{A(K_S \rightarrow \pi^+ \pi^-)} = \varepsilon + \varepsilon', \quad \frac{A(K_L \rightarrow \pi^0 \pi^0)}{A(K_S \rightarrow \pi^0 \pi^0)} = \varepsilon - 2\varepsilon'. \quad (5)$$

While $\varepsilon = (2.280 \pm 0.013) \times e^{i\frac{\pi}{4}} \times 10^{-3}$ parametrizes “indirect” CP violation originating from the fact that the mass eigenstates of the neutral K -meson system are not CP eigenstates, the quantity $\text{Re}(\varepsilon'/\varepsilon)$ measures “direct” CP violation in $K \rightarrow \pi\pi$ transitions. The CP-violating observable ε plays an important role to constrain the unitarity triangle⁷ and informs us in particular about a positive value of the Wolfenstein parameter η .

Despite enormous efforts, the experimental situation concerning $\text{Re}(\varepsilon'/\varepsilon)$ is still unclear at present. Whereas the CERN experiment NA31 finds $\text{Re}(\varepsilon'/\varepsilon) = (23 \pm 7) \times 10^{-4}$, which already indicates direct CP violation, the result $\text{Re}(\varepsilon'/\varepsilon) = (7.4 \pm 5.9) \times 10^{-4}$ of the Fermilab experiment E731 provides no unambiguous evidence for a non-zero effect. These results have been published already in 1993. In the near future, this unsatisfactory experimental situation will hopefully be clarified by improved measurements at CERN and Fermilab, as well as by the KLOE experiment at DAΦNE.

From a theoretical point of view, $\text{Re}(\varepsilon'/\varepsilon)$ is unfortunately not in much better shape. The corresponding calculations are very involved and suffer at present from large hadronic uncertainties⁷. Consequently, that observable will not allow a powerful test of the CP-violating sector of the Standard Model, unless the hadronic matrix elements of the relevant operators can be brought under better control. Probably the major goal of a possible future observation of $\text{Re}(\varepsilon'/\varepsilon) \neq 0$ would hence be the unambiguous exclusion of “super-weak” models of CP violation⁸.

In respect of testing the Standard Model description of CP violation, the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ are more promising. These decays – in particular the first one – are very clean from a theoretical point of view⁹. The branching ratios of these decays allow a determination of the unitarity triangle, if the top quark mass m_t and the CKM element $|V_{cb}|$ are used as an additional input. A detailed analysis shows that in particular $\sin(2\beta)$ can be extracted with respectable accuracy¹⁰. Thus, one has a powerful tool to probe physics beyond the Standard Model by comparing the value of $\sin(2\beta)$ determined in this way with the one extracted from CP violation in the “gold-plated” mode $B_d \rightarrow J/\psi K_S$ (see Subsection 3.1). Although the first experimental evidence for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ has recently been reported by the E787 collaboration¹¹, measurements of these rare kaon decays – in particular of $K_L \rightarrow \pi^0 \nu \bar{\nu}$ – are very challenging. Within the Standard Model, the expected branching ratios^{7,9} are $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (9.1 \pm 3.6) \times$

10^{-11} and $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = (2.8 \pm 1.7) \times 10^{-11}$. Nevertheless, there are plans to explore these important decays at BNL, FNAL and KEK.

To summarize the status of the kaon system, the observed CP violation in K -meson decays can be described successfully by the Standard Model at present. This feature is, however, not surprising, since so far only a single CP-violating observable, ε , has to be fitted. Consequently many different non-Standard Model descriptions of CP violation are imaginable³. From the brief discussion given above, it is obvious that the kaon system by itself cannot provide the whole picture of CP violation. Therefore it is essential to study CP violation outside this system. In this respect, the B system appears to be most promising, which is also reflected by the tremendous experimental efforts at future B -factory facilities. There are of course also other interesting systems to investigate CP violation and to search for signals of new physics, for instance the D -meson system¹², where sizeable mixing or CP-violating effects would signal new physics because of the tiny Standard Model “background”. In this presentation, I cannot discuss these systems in more detail and shall focus on B decays.

3. CP Violation in B Decays

The B -meson system is expected to provide a very fertile ground for testing the Standard Model description of CP violation. In this respect, the major role is played by non-leptonic B decays, which can be divided into three decay classes: decays receiving both tree and penguin contributions, pure tree decays, and pure penguin decays. A distinction is made between two types of penguin topologies: gluonic (QCD) and electroweak (EW) penguins related to strong and electroweak interactions, respectively. Because of the large top quark mass, also the latter operators play an important role in several processes¹³. In order to analyse non-leptonic B decays theoretically, one uses low-energy effective Hamiltonians, which are calculated by making use of the operator product expansion, yielding transition matrix elements of the following structure:

$$\langle f | \mathcal{H}_{\text{eff}} | i \rangle \propto \sum_k C_k(\mu) \langle f | Q_k(\mu) | i \rangle. \quad (6)$$

The operator product expansion allows us to separate the short-distance contributions to Eq. (6) from the long-distance contributions, which are described by perturbative Wilson coefficient functions $C_k(\mu)$ and non-perturbative hadronic matrix elements $\langle f | Q_k(\mu) | i \rangle$, respectively. As usual, μ denotes an appropriate renormalization scale.

In the case of $|\Delta B| = 1$, $\Delta C = \Delta U = 0$ transitions, which play the major role in the following discussion, we have

$$\mathcal{H}_{\text{eff}} = \mathcal{H}_{\text{eff}}(\Delta B = -1) + \mathcal{H}_{\text{eff}}(\Delta B = -1)^\dagger \quad (7)$$

with

$$\mathcal{H}_{\text{eff}}(\Delta B = -1) = \frac{G_F}{\sqrt{2}} \left[\sum_{j=u,c} V_{jq}^* V_{jb} \left\{ \sum_{k=1}^2 Q_k^{jq} C_k(\mu) + \sum_{k=3}^{10} Q_k^q C_k(\mu) \right\} \right]. \quad (8)$$

Here $\mu = \mathcal{O}(m_b)$, Q_k^{jq} are four-quark operators, and the label $q \in \{d, s\}$ corresponds to $b \rightarrow d$ and $b \rightarrow s$ transitions. The index k distinguishes between “current–current” ($k \in \{1, 2\}$), QCD ($k \in \{3, \dots, 6\}$) and electroweak ($k \in \{7, \dots, 10\}$) penguin operators, which are related to tree-level, QCD and electroweak penguin processes, respectively.

The evaluation of such low-energy effective Hamiltonians has been reviewed in Ref.¹⁴, where the four-quark operators are given explicitly and numerical values for their Wilson coefficient functions can be found.

3.1. CP Asymmetries in Neutral B Decays

A particularly simple and interesting situation arises, if we restrict ourselves to decays of neutral B_q mesons ($q \in \{d, s\}$) into CP self-conjugate final states $|f\rangle$ satisfying the relation $(\mathcal{CP})|f\rangle = \pm|f\rangle$. In that case the corresponding time-dependent CP asymmetry can be expressed as

$$a_{\text{CP}}(t) \equiv \frac{\Gamma(B_q^0(t) \rightarrow f) - \Gamma(\overline{B}_q^0(t) \rightarrow f)}{\Gamma(B_q^0(t) \rightarrow f) + \Gamma(\overline{B}_q^0(t) \rightarrow f)} = \mathcal{A}_{\text{CP}}^{\text{dir}}(B_q \rightarrow f) \cos(\Delta M_q t) + \mathcal{A}_{\text{CP}}^{\text{mix-ind}}(B_q \rightarrow f) \sin(\Delta M_q t), \quad (9)$$

where the direct CP-violating contributions have been separated from the mixing-induced CP-violating contributions, which are characterized by

$$\mathcal{A}_{\text{CP}}^{\text{dir}}(B_q \rightarrow f) \equiv \frac{1 - |\xi_f^{(q)}|^2}{1 + |\xi_f^{(q)}|^2} \quad \text{and} \quad \mathcal{A}_{\text{CP}}^{\text{mix-ind}}(B_q \rightarrow f) \equiv \frac{2 \text{Im} \xi_f^{(q)}}{1 + |\xi_f^{(q)}|^2}, \quad (10)$$

respectively. Here direct CP violation refers to CP-violating effects arising directly in the corresponding decay amplitudes, whereas mixing-induced CP violation is related to interference between B_q^0 - $\overline{B}_q^0$ mixing and decay processes. Note that the expression Eq. (9) has to be modified in the B_s case for $t \gtrsim 1/|\Delta\Gamma_s|$ because of the expected sizeable width difference $\Delta\Gamma_s$ ¹⁵.

In general, the observable

$$\xi_f^{(q)} \equiv e^{-i\phi_{\text{M}}^{(q)}} \frac{A(\overline{B}_q^0 \rightarrow f)}{A(B_q \rightarrow f)}, \quad \text{where} \quad \phi_{\text{M}}^{(q)} = \begin{cases} 2\beta & \text{for } q = d \\ 0 & \text{for } q = s \end{cases} \quad (11)$$

denotes the weak B_q^0 - $\overline{B}_q^0$ mixing phase, suffers from large hadronic uncertainties, which are introduced through the decay amplitudes A . There is, however, a very important special case, where these uncertainties cancel. It is given if $B_q \rightarrow f$ is dominated by a single CKM amplitude. In that case, $\xi_f^{(q)}$ takes the simple form

$$\xi_f^{(q)} = \mp \exp \left[-i \left(\phi_{\text{M}}^{(q)} - \phi_{\text{D}}^{(f)} \right) \right], \quad (12)$$

where $\phi_{\text{D}}^{(f)}$ is a characteristic weak decay phase, which is given by

$$\phi_{\text{D}}^{(f)} = \begin{cases} -2\gamma & \text{for dominant } \bar{b} \rightarrow \bar{u} u \bar{r} \text{ CKM amplitudes in } B_q \rightarrow f \\ 0 & \text{for dominant } \bar{b} \rightarrow \bar{c} c \bar{r} \text{ CKM amplitudes in } B_q \rightarrow f. \end{cases} \quad (13)$$

The label $r \in \{d, s\}$ distinguishes between $\bar{b} \rightarrow \bar{d}$ and $\bar{b} \rightarrow \bar{s}$ transitions.

Important applications and well-known examples of this formalism are the decays $B_d \rightarrow J/\psi K_S$ and $B_d \rightarrow \pi^+ \pi^-$. If one goes through the relevant Feynman diagrams contributing to the former channel (for a detailed recent discussion, see Ref.¹³), one finds that it is

dominated to excellent accuracy by the $\bar{b} \rightarrow \bar{c}c\bar{s}$ CKM amplitude. Therefore the decay phase vanishes and we have

$$\mathcal{A}_{\text{CP}}^{\text{mix-ind}}(B_d \rightarrow J/\psi K_S) = +\sin[-(2\beta - 0)]. \quad (14)$$

Since Eq. (12) applies to excellent accuracy to the decay $B_d \rightarrow J/\psi K_S$ – the point is that penguins enter essentially with the same weak phase as the leading tree contribution – it is usually referred to as the “gold-plated” mode to measure the CKM angle β ¹⁶.

In the case of $B_d \rightarrow \pi^+\pi^-$, mixing-induced CP violation would measure $-\sin(2\alpha)$ in a clean way through

$$\mathcal{A}_{\text{CP}}^{\text{mix-ind}}(B_d \rightarrow \pi^+\pi^-) = -\sin[-(2\beta + 2\gamma)] = -\sin(2\alpha), \quad (15)$$

if there were no penguin contributions present. However, such contributions are there and destroy the cleanliness of Eq. (15). The corresponding hadronic uncertainties were discussed by many authors in the literature. There are even methods to control these uncertainties in a quantitative way. Unfortunately, these strategies are usually rather challenging in practice. The most important examples are the $B \rightarrow \pi\pi$ isospin triangles proposed by Gronau and London¹⁷, and an approach using $B \rightarrow \rho\pi$ modes¹⁸. An approximate method to correct for the penguin uncertainties in $B_d \rightarrow \pi^+\pi^-$ that appears to be promising for the early days of the B -factory era was proposed in Ref.¹⁹. For a detailed discussion of these and other strategies, the reader is referred to reviews on this topic, for instance Refs.^{13,20}.

A decay appearing frequently as a tool to determine the CKM angle γ is $B_s \rightarrow \rho^0 K_S$. In that case, however, penguins are expected to lead to serious problems – more serious than in $B_d \rightarrow \pi^+\pi^-$ – so that this mode appears to be the “wrong” way to extract γ ¹³. Other strategies allowing meaningful determinations of this angle will be discussed below.

3.2. CP Violation in Non-leptonic Penguin Modes as a Probe of New Physics

In order to test the Standard Model description of CP violation, penguin-induced modes play an important role. Because of the loop-suppression of these “rare” processes, it is possible – and indeed it is the case in several specific model calculations – that new-physics contributions to these decays are of similar magnitude as those of the Standard Model³. An example is the $b \rightarrow d$ penguin mode $B_d \rightarrow K^0 \bar{K}^0$ (for an analysis of new-physics effects, see Ref.²¹). If one assumes that penguins with internal top quarks play the dominant role in this decay, the weak B_d^0 – $\bar{B}_d^0$ mixing and $B_d \rightarrow K^0 \bar{K}^0$ decay phases cancel in the corresponding observable $\xi_{K^0 \bar{K}^0}^{(d)}$, implying *vanishing* CP violation in that decay. Consequently one would conclude that a measurement of non-vanishing CP violation in $B_d \rightarrow K^0 \bar{K}^0$ would signal physics beyond the Standard Model. However, long-distance effects related to penguins with internal charm and up quarks may easily spoil the assumption of top quark dominance^{13,22}. These contributions may lead to sizeable CP violation in $B_d \rightarrow K^0 \bar{K}^0$ even within the Standard Model²³, so that a measurement of such CP asymmetries would not necessarily imply new physics, as claimed in several previous papers. Unfortunately a measurement of these effects will be difficult, since the Standard Model expectation for the corresponding branching ratio is $\mathcal{O}(10^{-6})$, which is still one order of magnitude below the recent CLEO bound²⁴ $\text{BR}(B_d \rightarrow K^0 \bar{K}^0) < 1.7 \times 10^{-5}$.

More promising in this respect and – more importantly – to search for physics beyond the Standard Model, is the $b \rightarrow s$ penguin mode $B_d \rightarrow \phi K_S$. The branching ratio for

this decay is expected to be of $\mathcal{O}(10^{-5})$ and may be large enough to investigate this channel at future B factories. Interestingly there is, to a very good approximation, no non-trivial CKM phase present in the corresponding decay amplitude¹³, so that direct CP violation vanishes and mixing-induced CP violation measures simply the weak $B_d^0\text{--}\overline{B}_d^0$ mixing phase. This statement does *not* require the questionable assumption of top quark dominance in penguin amplitudes. Consequently, an important probe for new physics in $b \rightarrow s$ flavour-changing neutral-current processes is provided by the relation

$$\mathcal{A}_{\text{CP}}^{\text{mix-ind}}(B_d \rightarrow J/\psi K_S) = \mathcal{A}_{\text{CP}}^{\text{mix-ind}}(B_d \rightarrow \phi K_S) = -\sin(2\beta), \quad (16)$$

which holds within the Standard Model framework. The theoretical accuracy of this relation is limited by certain neglected terms that are CKM-suppressed by $\mathcal{O}(\lambda^2)$ and may lead to tiny direct CP-violating asymmetries in $B_d \rightarrow \phi K_S$ of at most $\mathcal{O}(1\%)$ ¹³. The importance of $B_d \rightarrow \phi K_S$ and similar modes, such as $B_d \rightarrow \eta' K_S$, to search for new physics in $b \rightarrow s$ transitions has recently been emphasized by several authors^{13,25}.

3.3. The B_s System in the Light of $\Delta\Gamma_s$

In the B_s system very rapid $B_s^0\text{--}\overline{B}_s^0$ oscillations are expected, requiring an excellent vertex resolution system. Studies of CP violation in B_s decays are therefore regarded as being very difficult. An alternative route to investigate CP-violating effects may be provided by the width difference $\Delta\Gamma_s/\Gamma_s = \mathcal{O}(20\%)$ arising from CKM-favoured $b \rightarrow c\bar{c}s$ transitions into final states that are common to both B_s^0 and $\overline{B}_s^0$. Because of this width difference, already *untagged* data samples of B_s decays may exhibit CP-violating effects¹⁵.

In the recent literature^{26,27} (for a review, see for instance Ref.²⁸), several “untagged strategies” to extract the CKM angle γ have been proposed, using for example angular distributions in $B_s \rightarrow K^{*+}K^{*-}$, $K^{*0}\overline{K}^{*0}$ or $B_s \rightarrow D^*\phi$, $D_s^{*\pm}K^{*\mp}$ decays. To illustrate these strategies in more detail, let us focus on an approach to determine γ from untagged $B_s \rightarrow K^0\overline{K}^0$ and $B_s \rightarrow K^+K^-$ decays that was proposed in Ref.²⁶. Neglecting tiny EW penguin contributions and using the $SU(2)$ isospin symmetry of strong interactions, the corresponding untagged rates, which are defined by

$$\Gamma[f(t)] \equiv \Gamma(B_s^0(t) \rightarrow f) + \Gamma(\overline{B}_s^0(t) \rightarrow f), \quad (17)$$

take the form^{26,29}

$$\Gamma[K^0\overline{K}^0(t)] = \mathcal{C} |P'_s|^2 e^{-\Gamma_L^{(s)}t}, \quad \Gamma[K^+K^-(t)] = \mathcal{C} |P'_s|^2 \left[a e^{-\Gamma_L^{(s)}t} + b e^{-\Gamma_H^{(s)}t} \right], \quad (18)$$

where $\mathcal{C}$ is a trivially calculable phase-space factor and

$$a = 1 - 2r_s \cos \delta_s \cos \gamma + r_s^2 \cos^2 \gamma, \quad b = r_s^2 \sin^2 \gamma, \quad (19)$$

with $r_s \equiv |T'_s|/|P'_s|$, $\delta_s \equiv \delta_{T'_s} - \delta_{P'_s}$. The amplitudes P'_s and T'_s denote QCD penguin and colour-allowed $\bar{b} \rightarrow \bar{u}u\bar{s}$ current–current operator contributions, respectively. Since the untagged $B_s \rightarrow K^0\overline{K}^0$ rate measures $\mathcal{C} |P'_s|^2$, which gives the normalization of the untagged $B_s \rightarrow K^+K^-$ rate, both a and b can be determined. If the amplitude ratio r_s is known, the CKM angle γ and the strong phase δ_s can be extracted from these observables. Since $|P'_s|$ is measured through the untagged $B_s \rightarrow K^0\overline{K}^0$ rate, this required

input actually corresponds to $|T'_s|$, which can be fixed, e.g. through $B^+ \rightarrow \pi^+\pi^0$ and $SU(3)$ flavour symmetry, “factorization”, or hopefully lattice gauge theory one day. The required information about $|T'_s|$ introduces some model-dependence into the extracted value of γ . However, as we will see in more detail in Subsection 3.6, it is possible to derive interesting *bounds* on γ from the untagged $B_s \rightarrow K\bar{K}$ rates that do not suffer from such a model dependence²⁹.

Let me note that the B_s system provides also interesting probes for physics beyond the Standard Model. Important examples are the decays $B_s \rightarrow D_s^+ D_s^-$ and $B_s \rightarrow J/\psi\phi$. The latter is the counterpart of the “gold-plated” mode $B_d \rightarrow J/\psi K_S$ to measure β . These transitions are dominated by a single CKM amplitude and allow – in principle even from their untagged data samples²⁶ – the extraction of a CP-violating weak phase $\phi_{\text{CKM}} \equiv 2\lambda^2\eta$, which is expected to be of $\mathcal{O}(0.03)$ within the Standard Model⁷. Consequently, an extracted value of ϕ_{CKM} that is much larger than this Standard Model expectation would signal new-physics contributions to $B_s^0\text{--}\bar{B}_s^0$ mixing. The untagged $B_s \rightarrow K^0\bar{K}^0$ rate is also interesting to search for new physics. If future experiments should find that its time evolution depends – in contrast to Eq. (18) – on *two* exponentials, we would have an indication for physics beyond the Standard Model²⁹.

3.4. Extracting CKM Angles with Amplitude Relations

Since mixing effects are absent in the charged B system, the measurement of a non-vanishing CP asymmetry in a charged B decay would give us unambiguous evidence for direct CP violation, thereby ruling out “superweak” models. Such CP asymmetries arise from interference between decay amplitudes with both different CP-violating weak and CP-conserving strong phases. Whereas the weak phases are related to the CKM matrix, the strong phases are induced by strong final-state interaction effects and introduce severe theoretical uncertainties into the calculation, destroying in general the clean relation of the CP asymmetry to the phases of the CKM matrix.

Nevertheless, there are decays of charged B mesons that play an important role to extract angles of the unitarity triangle, in particular for γ . In this case amplitude relations – either exact or approximate ones based on flavour symmetries – are used. A recent review of these methods can be found in Ref.¹³. The “prototype” is the approach to determine γ with the help of triangle relations among the $B^\pm \rightarrow DK^\pm$ decay amplitudes proposed by Gronau and Wyler³⁰. Unfortunately the corresponding triangles are expected to be very “squashed” ones. Moreover one has to deal with additional experimental problems³¹, so that this approach is very difficult from a practical point of view. Recently more refined variants have been proposed by Atwood, Dunietz and Soni³¹.

About three years ago, several methods to extract CKM angles were presented by Gronau, Hernández, London and Rosner, who have combined the $SU(3)$ flavour symmetry of strong interactions with certain plausible dynamical assumptions to derive relations among $B \rightarrow \pi\pi, \pi K, K\bar{K}$ decay amplitudes³². This approach has been very popular over the recent years and requires only a measurement of the relevant branching ratios. A closer look shows, however, that it has – despite its attractiveness – theoretical limitations: the $SU(3)$ relations are not valid exactly, QCD penguins with internal charm and up quarks play an important role in several cases, and interestingly also EW penguins lead to complications. In order to eliminate the EW penguin contributions, usually very involved strategies are needed. A detailed discussion of all these methods (mainly to extract γ) is beyond the

scope of this presentation and the reader is referred to a recent review¹³ and references therein.

3.5. Constraining γ with $B_{u,d} \rightarrow \pi K$ Modes

A simple approach to determine γ with the help of the branching ratios for $B^+ \rightarrow \pi^+ K^0$, $B_d^0 \rightarrow \pi^- K^+$ and their charge-conjugates was proposed in Ref.³³ (see also Ref.¹³). Similarly as the $B_s \rightarrow K\bar{K}$ strategy²⁶ discussed in Subsection 3.3, it makes use of the fact that the general phase structure of the corresponding decay amplitudes is known reliably within the Standard Model. If the magnitude of the current-current amplitude T' contributing to $B_d^0 \rightarrow \pi^- K^+$ is known – it can be fixed similarly to $|T'_s|$, e.g. through $B^+ \rightarrow \pi^+ \pi^0$ and $SU(3)$ flavour symmetry, “factorization”, or hopefully lattice gauge theory one day – two amplitude triangles can be constructed, allowing in particular the extraction of γ . This approach is promising for future B -physics experiments, since it requires only time-independent measurements of branching ratios at the $\mathcal{O}(10^{-5})$ level. If one measures in addition the branching ratios for $B^+ \rightarrow \pi^0 K^+$ and its charge-conjugate, also the $b \rightarrow s$ electroweak penguin amplitude can be determined, which is – among other things – an interesting probe for new physics³⁴.

Recently the CLEO collaboration²⁴ has reported the first observation of the decays $B^+ \rightarrow \pi^+ K^0$ and $B_d^0 \rightarrow \pi^- K^+$. At present only combined branching ratios, i.e. averaged over decays and their charge-conjugates, are available with large experimental uncertainties. Therefore it is not yet possible to extract γ from the triangle construction proposed in Ref.³³. The recent CLEO measurements allow us, however, to derive *constraints* on γ , which are of the form

$$0^\circ \leq \gamma \leq \gamma_0 \quad \vee \quad 180^\circ - \gamma_0 \leq \gamma \leq 180^\circ \quad (20)$$

and are hence complementary to the presently allowed range

$$42^\circ \lesssim \gamma \lesssim 135^\circ \quad (21)$$

for that angle arising from the usual fits of the unitarity triangle⁷. This interesting feature has recently been pointed out in Ref.³⁵. The quantity γ_0 in Eq. (20) depends both on

$$R \equiv \frac{\text{BR}(B_d \rightarrow \pi^\mp K^\pm)}{\text{BR}(B^\pm \rightarrow \pi^\pm K)} = \frac{\text{BR}(B_d^0 \rightarrow \pi^- K^+) + \text{BR}(\bar{B}_d^0 \rightarrow \pi^+ K^-)}{\text{BR}(B^+ \rightarrow \pi^+ K^0) + \text{BR}(B^- \rightarrow \pi^- \bar{K}^0)}, \quad (22)$$

i.e. the ratio of combined branching ratios, and on the amplitude ratio $r \equiv |T'|/|P'|$ of the current-current and penguin operator contributions to $B_d \rightarrow \pi^\mp K^\pm$.

While the constraints on γ require knowledge about r for $R > 1$, bounds on γ can always be obtained *independently* of r , if R is found experimentally to be smaller than 1. The point is that γ_0 takes a maximal value

$$\gamma_0^{\text{max}} = \arccos(\sqrt{1 - R}), \quad (23)$$

which depends only on the ratio R of combined $B_{u,d} \rightarrow \pi K$ branching ratios³⁵.

Let us take as an example the central value 0.65 of the recent CLEO measurements²⁴, yielding $R = 0.65 \pm 0.40$. This value corresponds to $\gamma_0^{\text{max}} = 54^\circ$ and implies the range $0^\circ \leq \gamma \leq 54^\circ \vee 126^\circ \leq \gamma \leq 180^\circ$, which has only the small overlap $42^\circ \lesssim \gamma \leq 54^\circ \vee$

$126^\circ \leq \gamma \lesssim 135^\circ$ with the range (21). The two pieces of this range are distinguished by the sign of $\cos \delta$, where δ is the CP-conserving strong phase shift between the T' and P' amplitudes. Using arguments based on “factorization”, one expects $\cos \delta > 0$ corresponding to the former interval of that range³⁵ (for a recent model calculation, see Ref.³⁶). Consequently, once more data come in confirming $R < 1$, the decays $B_d \rightarrow \pi^\mp K^\pm$ and $B^\pm \rightarrow \pi^\pm K$ may put the Standard Model to an interesting test and could open a window to new physics. Effects of physics beyond the Standard Model in $B_{u,d} \rightarrow \pi K$ decay amplitudes have been analysed in a recent paper³⁷. A detailed study of various implications of the bounds on γ discussed above has recently been performed in Ref.³⁸, where the issue of new physics has also been addressed.

3.6. Towards Extractions of γ from $B_{u,d} \rightarrow \pi K$ and Untagged $B_s \rightarrow K\bar{K}$ Decays

At first sight, the observables a and b specified in Eq. (19), which can be extracted from the time evolutions of the untagged $B_s \rightarrow K^0\bar{K}^0$ and $B_s \rightarrow K^+K^-$ decay rates, provide a similar bound on γ as the combined $B_{u,d} \rightarrow \pi K$ rates. The point is that $R_s \equiv a + b$ corresponds exactly to the ratio R of combined $B_{u,d} \rightarrow \pi K$ branching ratios.

A closer look shows, however, that it is possible to derive a more elaborate bound on γ from the untagged $B_s \rightarrow K\bar{K}$ decay rates²⁹:

$$\frac{|1 - \sqrt{a}|}{\sqrt{b}} \leq |\cot \gamma| \leq \frac{1 + \sqrt{a}}{\sqrt{b}}, \quad (24)$$

which corresponds to the allowed range

$$\gamma_1 \leq \gamma \leq \gamma_2 \quad \vee \quad 180^\circ - \gamma_2 \leq \gamma \leq 180^\circ - \gamma_1 \quad (25)$$

with

$$\gamma_1 = \operatorname{arccot} \left(\frac{1 + \sqrt{a}}{\sqrt{b}} \right), \quad \gamma_2 = \operatorname{arccot} \left(\frac{|1 - \sqrt{a}|}{\sqrt{b}} \right). \quad (26)$$

It can be shown²⁹ that the bound Eq. (25) always excludes a range around $\gamma = 90^\circ$ larger than the bound $|\cos \gamma| \geq \sqrt{1 - R_s}$ arising for $R_s < 1$, provided $R_s \neq \sqrt{a}$. For $R_s = \sqrt{a}$, both bounds exclude the same region around $\gamma = 90^\circ$.

So far, we have discussed the $B_{u,d} \rightarrow \pi K$ and $B_s \rightarrow K\bar{K}$ decays separately. As we have just seen, interesting and stringent bounds on γ may arise from the corresponding observables. The goal is, however, not only to constrain γ , but to *determine* this angle eventually. If we consider $B_{u,d} \rightarrow \pi K$ and $B_s \rightarrow K\bar{K}$ separately, information about colour-allowed $\bar{b} \rightarrow \bar{u}u\bar{s}$ current-current amplitudes is needed to accomplish this task^{26,33}, which introduces hadronic uncertainties into the extracted values of γ . Such an input can be avoided by relating r_s and r straightforwardly through

$$\left(\frac{r_s}{r} \right)^2 \equiv \zeta, \quad (27)$$

where ζ parametrizes $SU(3)$ breaking related to the s and u, d spectator quarks in the modes $B_s \rightarrow K\bar{K}$ and $B_{u,d} \rightarrow \pi K$, respectively. As has recently been pointed out in Ref.²⁹, the observables of the $B_{u,d} \rightarrow \pi K$ and $B_s \rightarrow K\bar{K}$ modes allow a simultaneous determination of γ and r, δ, r_s, δ_s (up to certain discrete ambiguities) as a function of the $SU(3)$ breaking parameter ζ . Keeping ζ explicitly in the formulae presented in Ref.²⁹ will

turn out to be useful once $SU(3)$ breaking can be controlled in a quantitative way. As a first “guess”, we may use $\zeta = 1$, which appears to be a rather mild $SU(3)$ assumption, since ζ describes $SU(3)$ breaking originating only from different spectator quarks, which should play a minor role for the decay dynamics. One may also vary ζ within a reasonable range to get some feeling for the uncertainties arising from possible $SU(3)$ breaking.

It is interesting to note that the approach sketched above works also for vanishing strong phases, and for a particular scenario of new physics, where the $B_s \rightarrow K\bar{K}$ and $B_{u,d} \rightarrow \pi K$ decay amplitudes are dominated by the Standard Model diagrams and $B_s^0\text{--}\bar{B}_s^0$ mixing receives CP-violating new-physics contributions. The observables of untagged $B_s \rightarrow K^0\bar{K}^0$ and $B_s \rightarrow D_s^+ D_s^-$ decays provide an interesting test of whether such a scenario of physics beyond the Standard Model is actually realized in nature²⁹.

4. Concluding Remarks and Outlook

In conclusion, we have seen that the kaon system – the only one where CP violation has been observed to date – cannot provide the whole picture of that phenomenon. In addition to other interesting systems, for instance the D system, non-leptonic B -meson decays, where large CP asymmetries are expected within the Standard Model, are extremely promising to test the CKM picture of CP violation. More advanced experimental studies of CP-violating effects in the kaon system and the exploration of CP violation at B physics facilities are just ahead of us. In the foreseeable future, these experiments may bring unexpected results that could shed light on the physics beyond the Standard Model. Certainly an exciting future lies ahead of us!

Acknowledgements

I am very grateful to Michael Beyer and Thomas Mannel for inviting me to such an interesting and well-organized workshop, and for providing generous travel support.

References

1. N. Cabibbo, *Phys. Rev. Lett.* **10** (1963) 531; M. Kobayashi and T. Maskawa, *Progr. Theor. Phys.* **49** (1973) 652.
2. C. Jarlskog, *Phys. Rev. Lett.* **55** (1985) 1039; *Z. Phys.* **C29** (1985) 491.
3. For reviews, see for instance Y. Grossman, Y. Nir and R. Rattazzi, SLAC-PUB-7379 (1997) [hep-ph/9701231]; M. Gronau and D. London, *Phys. Rev.* **D55** (1997) 2845; Y. Nir and H.R. Quinn, *Annu. Rev. Nucl. Part. Sci.* **42** (1992) 211; R. Fleischer, CERN-TH/97-241 (1997) [hep-ph/9709291], invited plenary talk given at the 7th International Symposium on Heavy Flavor Physics, Santa Barbara, California, 7-11 July 1997, to appear in the proceedings.
4. L. Wolfenstein, *Phys. Rev. Lett.* **51** (1983) 1945.
5. L.L. Chau and W.-Y. Keung, *Phys. Rev. Lett.* **53** (1984) 1802; C. Jarlskog and R. Stora, *Phys. Lett.* **B208** (1988) 268.
6. J.H. Christenson, J.W. Cronin, V.L. Fitch and R. Turlay, *Phys. Rev. Lett.* **13** (1964) 138.
7. For a review, see A.J. Buras and R. Fleischer, University of Karlsruhe preprint TTP97-15 (1997) [hep-ph/9704376].

8. L. Wolfenstein, *Phys. Rev. Lett.* **13** (1964) 562.
9. G. Buchalla and A.J. Buras, *Nucl. Phys.* **B412** (1994) 106 and **B400** (1993) 225.
10. G. Buchalla and A.J. Buras, *Phys. Rev.* **D54** (1996) 6782; *Phys. Lett.* **B333** (1994) 221; A.J. Buras, *Phys. Lett.* **B333** (1994) 476.
11. S. Adler et al., E787 collaboration, *Phys. Rev. Lett.* **79** (1997) 2204.
12. For a review, see E. Golowich, [hep-ph/9706548].
13. For a review, see R. Fleischer, *Int. J. Mod. Phys.* **A12** (1997) 2459.
14. G. Buchalla, A.J. Buras and M.E. Lautenbacher, *Rev. Mod. Phys.* **68** (1996) 1125.
15. I. Dunietz, *Phys. Rev.* **D52** (1995) 3048.
16. A.B. Carter and A.I. Sanda, *Phys. Rev. Lett.* **45** (1980) 952; *Phys. Rev.* **D23** (1981) 1567; I.I. Bigi and A.I. Sanda, *Nucl. Phys.* **B193** (1981) 85.
17. M. Gronau and D. London, *Phys. Rev. Lett.* **65** (1990) 3381.
18. H. Lipkin, Y. Nir, H. Quinn and A. Snyder, *Phys. Rev.* **D44** (1991) 1454; A. Snyder and H. Quinn, *Phys. Rev.* **D48** (1993) 2139.
19. R. Fleischer and T. Mannel, *Phys. Lett.* **B397** (1997) 269.
20. For reviews, see for instance Y. Nir and H.R. Quinn, in *B Decays*, ed. S. Stone (World Scientific, Singapore, 1994), p. 362; I. Dunietz, *ibid.*, p. 393; H.R. Quinn, *Nucl. Phys.* **B** (Proc. Suppl.) **50** (1996) 17; R. Fleischer, Ref.¹³.
21. M.P. Worah, *Phys. Rev.* **D54** (1996) 2198.
22. A.J. Buras and R. Fleischer, *Phys. Lett.* **B341** (1995) 379. Recently this point has also been raised by M. Ciuchini et al., CERN-TH/97-030 (1997) [hep-ph/9703353].
23. R. Fleischer, *Phys. Lett.* **B341** (1994) 205.
24. J.P. Alexander, invited plenary talk given at the 2nd International Conference on *B* Physics and CP Violation, Honolulu, Hawaii, 24-27 March 1997, to appear in the proceedings; F. Würthwein, CALT-68-2121 (1997) [hep-ex/9706010].
25. Y. Grossman and M.P. Worah, *Phys. Lett.* **B395** (1997) 241; M. Ciuchini et al., *Phys. Rev. Lett.* **79** (1997) 978; D. London and A. Soni, *Phys. Lett.* **B407** (1997) 61.
26. R. Fleischer and I. Dunietz, *Phys. Rev.* **D55** (1997) 259.
27. R. Fleischer and I. Dunietz, *Phys. Lett.* **B387** (1996) 361.
28. R. Fleischer, University of Karlsruhe preprint TTP97-21 (1997) [hep-ph/9705404], invited plenary talk given at the 2nd International Conference on *B* Physics and CP Violation, Honolulu, Hawaii, 24-27 March 1997, to appear in the proceedings.
29. R. Fleischer, CERN-TH/97-281 (1997) [hep-ph/9710331].
30. M. Gronau and D. Wyler, *Phys. Lett.* **B265** (1991) 172.
31. D. Atwood, I. Dunietz and A. Soni, *Phys. Rev. Lett.* **78** (1997) 3257.
32. M. Gronau et al., *Phys. Rev.* **D52** (1995) 6374, and references therein.
33. R. Fleischer, *Phys. Lett.* **B365** (1996) 399.
34. A.J. Buras and R. Fleischer, *Phys. Lett.* **B365** (1996) 390.
35. R. Fleischer and T. Mannel, University of Karlsruhe preprint TTP97-17 (1997) [hep-ph/9704423].
36. A. Ali and C. Greub, DESY-97-126 (1997) [hep-ph/9707251].
37. R. Fleischer and T. Mannel, University of Karlsruhe preprint TTP97-22 (1997) [hep-ph/9706261].
38. Y. Grossman, Y. Nir, S. Plaszczynski and M. Schune, SLAC-PUB-7622 (1997) [hep-ph/9709288].